

Mansonella perstans

Mansonella perstans	3
----------------------------------	----------

Mansonella perstans

Mansonella perstans (formerly *Dipetalonema perstans*) is a nematode transmitted by *Culicoides* insects. These midges have an aggressive and very annoying biting behaviour principally at dusk but also to a lesser extent at night and during the day. They bite predominantly on hands, arms and head. They often fly in swarms around the face. In view of their large numbers, they can form a very severe plague. Control is difficult.




© CD-ROM ILLUSTRATED LECTURE NOTES ON TROPICAL MEDICINE

Midges. *Culicoides* sp. Vectors of *Mansonella filaria*. Copyright ITM

Mansonella perstans

© CD-ROM ILLUSTRATED LECTURE NOTES ON TROPICAL MEDICINE



Map showing areas endemic for *Mansonella perstans* filariosis. Copyright ITM

The infection is widely distributed in Africa but is more localised in Central and South America. *M. perstans* does not occur in Asia.

The adult worms live in body cavities (peritoneum, pleura and pericardium) and in perirenal fat. Most patients infected with *Mansonella perstans* are asymptomatic. A number of different symptoms and allergic reactions are sometimes ascribed to this parasite, but the disease spectrum has not yet been fully established.

The diagnosis is established by detecting the typical small microfilariae in the peripheral blood.

If asymptomatic, no treatment is necessary. A 6-week course of doxycycline with 200 mg/day resulted in a 97% cure-rate at 12 months in a study in Mali. Ivermectin, albendazole and DEC are inactive.

LAST UPDATED BY ADMIN ON JULY 15TH, 2022